

Drug Information Sheet("Kusuri-no-Shiori")

Injection

Published: 11/2012

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Kindaly 3D

Active ingredient:Powder Dialysate for Hemodialysis

Dosage form:Dialysate

Print on wrapping:



Effects of this medicine

This medicine purifies blood excreting waste matter in the blood by diffusion/osmosis phenomenon through dialysis membrane of the dialyzer.

It also corrects electrolyte balance and extra fluid balance.

It is usually used as a perfusate of artificial kidney on dialysis therapy for chronic renal failure with hypercalcemia.

There are varieties of compositions on dialysate, this medicine is used when high bicarbonate concentration dialysate is likely to cause excessive alkalosis, when glycemic control is difficult by low sugar concentration dialysate or when dialysate high in potassium or magnesium fails to adequately improve hyperkalemia or hypermagnesemia.

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you have liver disorder and diabetes mellitus.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, for adults, it is used as a perfusate to undergo dialysis.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include leg cramp. Adverse reactions reported as dialysis therapy include hypotension, shock, increased blood pressure, convulsion, feeling ill, bone related complication (fracture), ectopic calcinosis (pain, itch), hyperglycemia, weight gain and dialysisdisequilibrium syndrome (headache, nausea/vomiting, consciousness disturbance, malaise).

If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.