



# Drug Information Sheet("Kusuri-no-Shiori")

Injection

Revised: 02/2026

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**Carbazochrome Sodium Sulfonate Injection 50mg  
"FUSO"

**Active ingredient:**Carbazochrome sodium sulfonate hydrate

**Dosage form:**injection

**Imprint or print on wrapping:**



## Effects of this medicine

This medicine reinforces blood vessels to enhance resistance and prevents bleeding from the vessels. It is usually used to treat bleeding tendency (purpura, etc.) caused by decreased microvascular resistance, bleeding from skin or mucosa and inner membrane, fundal/renal/uterine bleeding, and abnormal bleeding during/after surgery.

## The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods. If you have hereditary fructose intolerance.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, inject this medicine intravenously or by intravenous infusion. Ask the doctor about the treatment span.

## Precautions while taking this medicine

- This medicine may cause orange-yellow discoloration of urine.
- This medicine may affect laboratory values. When you take clinical laboratory tests, make sure to tell your doctor that you are using this medicine.

## Possible adverse reactions to this medicine

The most commonly reported adverse reaction includes rash. If the symptom occurs, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- facial pallor, cold sweat, dizziness on standing up [shock, anaphylaxis]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.